

ON A CLASS OF NEARLY BINARY MATROIDS

JAMES OXLEY AND JESSE TAYLOR

ABSTRACT. We give an excluded-minor characterization for the class of matroids M in which $M \setminus e$ or M/e is binary for all e in $E(M)$. This class is closely related to the class of matroids in which every member is binary or can be obtained from a binary matroid by relaxing a circuit-hyperplane. We also provide an excluded-minor characterization for the second class.

1. INTRODUCTION

In this paper, we consider a class of matroids that are close to being binary. Specifically, we consider the class $\mathcal{Z}$ of matroids M such that $M \setminus e$ or M/e is binary for all e in $E(M)$. The main result is an excluded-minor characterization of $\mathcal{Z}$. Except where otherwise noted the notation and terminology follow [8].

It is well known that if H is a circuit and a hyperplane of a matroid M , then there is another matroid M' on $E(M)$ whose bases are the bases of M together with H . We say that M' is obtained from M by *relaxing* the *circuit-hyperplane* H and call M' a *relaxation* of M . A class of matroids that arises naturally in determining the excluded minors for $\mathcal{Z}$ is $\mathcal{R}$, those matroids M such that M is binary or M is a relaxation of a binary matroid.

The rank-three whirl is denoted by $\mathcal{W}^3$, while P_6 is the six-element rank-three matroid that has a single triangle as its only non-spanning circuit. Let Q_6 and R_6 be the six-element matroids of rank three for which geometric representations are given in Figure 1. Evidently $R_6 \cong U_{2,4} \oplus_2 U_{2,4}$. Let K be the seven-element rank-two matroid that is obtained by adding elements in parallel to three of the elements of $U_{2,4}$. The matroid K is depicted with its dual in Figure 2. In Section 2, we prove that $\mathcal{Z}$ and $\mathcal{R}$ are minor-closed classes of matroids and establish some excluded minors of each. We also introduce Cunningham and Edmonds's canonical tree decomposition of a 2-connected matroid, along with some preliminaries.

Let N be a matroid. A matroid M is *N -fragile* if, for each element e of $E(M)$, at least one of $M \setminus e$ and M/e has no N -minor (see, for example, [4]). The class of N -fragile matroids is clearly minor-closed. The main result of this paper determines the set of excluded minors for the class of $U_{2,4}$ -fragile

Date: March 3, 2013.

1991 *Mathematics Subject Classification.* 05B35.

The authors were supported by the National Security Agency and by an NSF VIGRE Grant, respectively.

matroids. Section 3 is devoted to proving this result and another related result, both of which are stated below.




FIGURE 1. Geometric representations of the six-element rank-three matroids Q_6 and R_6 .




FIGURE 2. Representations of the matroid K and its (rank-5) dual K^* .

Theorem 1.1. *The set of excluded minors for the class of matroids M such that $M \setminus e$ or M/e is binary for all e in $E(M)$ is $\{Q_6, P_6, U_{3,6}, R_6, U_{2,4} \oplus U_{1,1}, U_{2,4} \oplus U_{0,1}\}$ together with all matroids that are obtained from connected binary matroids by relaxing two disjoint circuit-hyperplanes that partition the ground set.*

Theorem 1.2. *The set of excluded minors for the class of matroids in which every member is binary or can be obtained from a binary matroid by relaxing a circuit-hyperplane is $\{U_{2,5}, U_{3,5}, K, K^*, R_6, U_{2,4} \oplus U_{1,1}, U_{2,4} \oplus U_{0,1}\}$ together with all matroids that are obtained from connected binary matroids by relaxing two disjoint circuit-hyperplanes that partition the ground set.*

Let $\mathcal{D}$ denote the collection of all matroids that are obtained from connected binary matroids by relaxing two disjoint circuit-hyperplanes that partition the ground set. For an even integer r exceeding two, let M_r be the rank- r tipless binary spike, that is, the vector matroid of the binary matrix $[I_r | J_r - I_r]$ where J_r is the matrix of all ones. Labeling the columns of this matrix $e_1, e_2, \dots, e_{2r}$ in order, we see that $\{e_2, e_3, \dots, e_r, e_{r+1}\}$ and its complement are both circuit-hyperplanes of M_r . By relaxing these circuit-hyperplanes, we obtain a member of $\mathcal{D}$. Thus the sets of excluded minors in Theorems 1.1 and 1.2 are both infinite. However, these doubly relaxed

spikes are not the only members of $\mathcal{D}$. In Section 4, we will further discuss the complexity of $\mathcal{D}$.

2. PRELIMINARIES

This section will first note that both $\mathcal{Z}$ and $\mathcal{R}$ are minor-closed, and will then determine some excluded minors for each class.

Lemma 2.1. *The classes $\mathcal{Z}$ and $\mathcal{R}$ are both closed under duality and the taking of minors.*

This lemma is a straightforward consequence of the following result of Kahn [3] (see also [8, p. 115]).

Lemma 2.2. *Let X be a circuit-hyperplane of a matroid M , and let M' be the matroid obtained from M by relaxing X . Then $(M')^*$ is obtained from M^* by relaxing the circuit-hyperplane $E(M) - X$ of the latter. Moreover, when $e \in E(M) - X$ and $f \in X$,*

- (i) $M/e = M'/e$ and, unless M has e as a coloop, $M' \setminus e$ is obtained from $M \setminus e$ by relaxing the circuit-hyperplane X of the latter; and
- (ii) $M \setminus f = M' \setminus f$ and, unless M has f as a loop, M'/f is obtained from M/f by relaxing the circuit-hyperplane $X - f$ of the latter.

It is not difficult to deduce from the following result of Oxley [7] that the class $\mathcal{R}$ is contained in the class $\mathcal{Z}$. We say a matroid N is a *series extension* of a matroid M if $M = N/T$ and every element of T is in series with some element of M . We call N a *parallel extension* of M if N^* is a series extension of M^* . Note that this differs from the terminology used in [8].

Theorem 2.3. *The following two statements are equivalent for a matroid M .*

- (i) M is non-binary and, for every element e , at least one of $M \setminus e$ and M/e is binary.
- (ii) (a) Both $r(M)$ and $r^*(M)$ exceed two and M can be obtained from a connected binary matroid by relaxing a circuit-hyperplane; or
 - (b) M is isomorphic to a parallel extension of $U_{2,n}$ for some $n \geq 5$; or
 - (c) M is isomorphic to a series extension of $U_{n-2,n}$ for some $n \geq 5$; or
 - (d) M can be obtained from $U_{2,4}$ by series extension of a subset S of $E(U_{2,4})$ and parallel extension of a disjoint subset T of $E(U_{2,4})$ where S or T may be empty.

Some excluded minors for $\mathcal{Z}$ and $\mathcal{R}$ are easy to identify. We omit the routine argument that establishes the following.

Lemma 2.4. *The matroids $U_{2,4} \oplus U_{1,1}$, $U_{2,4} \oplus U_{0,1}$, and R_6 are excluded minors for both $\mathcal{Z}$ and $\mathcal{R}$.*

The following result from [3] will also be useful in identifying some excluded minors.

Lemma 2.5. *Let M' be obtained from M by relaxing the circuit-hyperplane X .*

- (i) *If M is connected, then M' is non-binary; and*
- (ii) *if M is n -connected, then so is M' .*

Let $\mathcal{D}$ denote the collection of all matroids that are obtained from connected binary matroids by relaxing two disjoint circuit-hyperplanes that partition the ground set.

Lemma 2.6. *All matroids in the collection $\mathcal{D}$ are in both $EX(\mathcal{Z})$ and $EX(\mathcal{R})$.*

Proof. Take a matroid M_2 in $\mathcal{D}$. Then M_2 can be obtained from a connected binary matroid M by relaxing two disjoint circuit-hyperplanes X and Y of M that partition $E(M)$. Note that because X and Y partition $E(M)$ and $E(M)$ is connected, M cannot have any circuits or cocircuits with fewer than four elements. Let M_X and M_Y denote the matroids obtained from M by relaxing X and Y , respectively, and take e in $E(M_2)$.

Suppose $e \in X$. By Lemma 2.2, M_2/e is obtained from M_Y/e by relaxing the circuit-hyperplane $X - e$ of the latter, and $M_Y/e = M/e$. If M/e is connected, then M_2/e is non-binary by Lemma 2.5. Now assume M/e is disconnected. Then M/e has at least two non-empty components, C_1 and C_2 . As Y is a spanning circuit in M/e , the last matroid has at least one component of rank zero, that is, M/e has a loop. But this contradicts the fact that all circuits of M have size at least four. We conclude that M_2/e is non-binary.

By Lemma 2.2 again, $M_2 \setminus e = M_Y \setminus e$, and this matroid is obtained from $M \setminus e$ by relaxing the circuit-hyperplane Y of the latter. Now $M \setminus e$ is connected by the dual of the argument above. Hence $M_2 \setminus e$ is non-binary by Lemma 2.5. The same arguments work for $f \in Y$ by replacing M_Y with M_X . Thus M_2 is not in $\mathcal{Z}$ and hence is not in $\mathcal{R}$.

As we saw above, any deletion $M_2 \setminus z$ equals $M_Y \setminus z$ or $M_X \setminus z$, depending on whether z is in X or Y . Both $M_Y \setminus z$, when $z \in X$, and $M_X \setminus z$, when $z \in Y$, can be obtained by relaxing a circuit-hyperplane in a binary matroid. By duality, any single-element contraction of M_2 can be obtained by relaxing a circuit-hyperplane in a binary matroid. Therefore any minor of M_2 is in $\mathcal{R}$, and so is in $\mathcal{Z}$. Thus M_2 is in $EX(\mathcal{R})$ and in $EX(\mathcal{Z})$. $\square$

The next two lemmas list some matroids that are excluded minors for exactly one of $\mathcal{R}$ and $\mathcal{Z}$. Their routine proofs are omitted.

Lemma 2.7. *The matroids $U_{2,5}$, $U_{3,5}$, K , and K^* are excluded minors for the class $\mathcal{R}$.*

Lemma 2.8. *The matroids Q_6 , P_6 , and $U_{3,6}$ are excluded minors for the class $\mathcal{Z}$.*

A class $\mathcal{N}$ of matroids is *1-rounded* [11] if every member of $\mathcal{N}$ is connected and, whenever e is an element of a connected matroid M having an $\mathcal{N}$ -minor, M has an $\mathcal{N}$ -minor using e .

Next we introduce Cunningham and Edmonds's tree decomposition for connected matroids that are not 3-connected [2]. Our treatment of this material will follow [8, pp. 307–310]. A *matroid-labeled tree* is a tree T with vertex set $\{M_1, M_2, \dots, M_k\}$ for some positive integer k such that

- (i) each M_i is a matroid;
- (ii) if M_{j_1} and M_{j_2} are joined by an edge e_i of T , then $E(M_{j_1}) \cap E(M_{j_2}) = \{e_i\}$, and $\{e_i\}$ is not a separator of M_{j_1} or M_{j_2} ; and
- (iii) if M_{j_1} and M_{j_2} are non-adjacent, then $E(M_{j_1}) \cap E(M_{j_2})$ is empty.

Let e be an edge of a matroid-labeled tree T and suppose e joins vertices labeled by M_1 and M_2 . Suppose that we contract e and relabel by $M_1 \oplus_2 M_2$ the composite vertex that results by identifying the endpoints of e . Then we retain a matroid-labeled tree and we denote this tree by T/e . This process can be repeated and since the operation of 2-sum is associative, for every subset $\{e_{i_1}, e_{i_2}, \dots, e_{i_m}\}$ of $E(T)$, the matroid-labeled tree $T/e_{i_1}, e_{i_2}, \dots, e_{i_m}$ is well-defined.

A *tree decomposition* of a 2-connected matroid M is a matroid-labeled tree T such that if $V(T) = \{M_1, M_2, \dots, M_k\}$ and $E(T) = \{e_1, e_2, \dots, e_{k-1}\}$, then

- (i) $E(M) = (E(M_1) \cup E(M_2) \cup \dots \cup E(M_k)) - \{e_1, e_2, \dots, e_{k-1}\}$;
- (ii) $|E(M_i)| \geq 3$ for all i unless $|E(M)| < 3$, in which case $k=1$ and $M_1=M$; and
- (iii) M is the matroid that labels the single vertex of $T/e_1, e_2, \dots, e_{k-1}$.

In general, a tree decomposition of a matroid is not unique. However, Cunningham and Edmonds were able to guarantee uniqueness of the *canonical tree decomposition* described in the following theorem from [2].

Theorem 2.9. *Let M be a 2-connected matroid. Then M has a tree decomposition T in which every vertex is labeled by a 3-connected matroid, a circuit, or a cocircuit, and there are no two adjacent vertices that are both labeled by circuits or are both labeled by cocircuits. Moreover, T is unique to within a relabeling of its edges.*

The canonical tree decomposition provides a unique way to break up a 2-connected matroid M into 3-connected pieces, circuits, and cocircuits. Moreover, we can reconstruct M from these pieces using the 2-sum operation with the common elements between matroids as basepoints. A basic property of the 2-sum operation is that M_1 and M_2 are isomorphic to minors of $M_1 \oplus_2 M_2$. The following result is well known; its routine proof is omitted.

Lemma 2.10. *Let M_1 and M_2 label vertices in a tree decomposition T of a connected matroid M . Let P be the path in T joining M_1 and M_2 and let p_1 and p_2 be the edges of P meeting M_1 and M_2 respectively. Then M has as a minor the 2-sum of M_1 and M_2 with respect to the basepoints p_1 and p_2 .*

3. MAIN RESULT

In this section we prove the main results of the paper, Theorem 1.1 and Theorem 1.2. We begin by finding all the disconnected excluded minors of each class. Due to the similarity of the proofs for each class, we will combine the arguments where possible.

Lemma 3.1. *Suppose $\mathcal{U} \in \{\mathcal{Z}, \mathcal{R}\}$. The only disconnected members of $EX(\mathcal{U})$ are $U_{2,4} \oplus U_{1,1}$ and $U_{2,4} \oplus U_{0,1}$.*

Proof. By Lemma 2.4, both $U_{2,4} \oplus U_{1,1}$ and $U_{2,4} \oplus U_{0,1}$ are in $EX(\mathcal{U})$. Now let M be an arbitrary disconnected member of $EX(\mathcal{U})$. As M is non-binary and disconnected, it has distinct components M_1 and M_2 where M_1 is non-binary. Since M_1 has a $U_{2,4}$ -minor and M_2 has a $U_{0,1}$ - or $U_{1,1}$ -minor, the lemma follows. $\square$

The following result [6] will be useful in our proofs.

Theorem 3.2. *Let M be a 3-connected matroid having rank and corank exceeding two.*

- (i) *If M is binary, then M has an $M(K_4)$ -minor.*
- (ii) *If M is non-binary, then M has one of $\mathcal{W}^3$, Q_6 , P_6 , and $U_{3,6}$ as a minor.*

Let B be a basis of a matroid M . For $e \in E(M) - B$, we denote by $C_M(e, B)$ the *fundamental circuit of e with respect to B* , that is, the unique circuit contained in $B \cup e$. Before finding the complete list of 2-connected excluded minors, we will need the following lemmas. The first lemma comes from [8, Section 1.5, Exercise 14]; its proof is omitted.

Lemma 3.3. *The following statements are equivalent for a matroid M .*

- (a) *M is a relaxation of some matroid.*
- (b) *M has a basis B such that $C_M(e, B) = B \cup e$ for every e in $E(M) - B$ and neither B nor $E(M) - B$ is empty.*

Lemma 3.4. *Let M be a matroid that can be obtained from a binary matroid N by relaxing a circuit-hyperplane X of the latter. If M contains a $\mathcal{W}^k$ -minor for some $k \geq 3$, then, in every $\mathcal{W}^k$ -minor of M , the rim elements are contained in X and no element of X is a spoke.*

Proof. Let M_1 be a $\mathcal{W}^k$ -minor of M . If e is in the rim of M_1 , then M_1/e is non-binary. But, for all f in $E(M) - X$, by Lemma 2.2, M/f is binary. Therefore $e \notin E(M) - X$, so $e \in X$. Now let s be a spoke of M_1 and t be

an adjacent rim element. Then $M_1/t \setminus s \cong \mathcal{W}^{k-1}$. Thus $M_1 \setminus s$ is non-binary, so $s \notin X$. $\square$

Lemma 3.5. *Let M be a connected non-binary matroid. Either M has an R_6 -, $U_{2,4} \oplus U_{0,1}$ -, or $U_{2,4} \oplus U_{1,1}$ -minor, or M is obtained from a 3-connected non-binary matroid M_0 by parallel extension of some subset T of $E(M_0)$ and series extension of a disjoint subset S of $E(M_0)$, where both S and T are possibly empty.*

Proof. Consider the canonical tree decomposition T of M . As M is non-binary, there must be a non-binary matroid M_0 labeling a vertex in the tree decomposition. Assume there is another vertex labeled by a non-binary matroid M_1 . Then, by Lemma 2.10, we see that M has an $M_0 \oplus_2 M_1$ -minor. Let p_1 be the basepoint of this 2-sum. Each of M_0 and M_1 is a connected non-binary matroid. Thus each has $U_{2,4}$ as a minor. Since $\{U_{2,4}\}$ is 1-rounded [1], each of M_0 and M_1 has a $U_{2,4}$ -minor using p_1 . Thus M has a $U_{2,4} \oplus_2 U_{2,4}$ -minor. But $U_{2,4} \oplus_2 U_{2,4} \cong R_6$, so the lemma holds when M_1 exists.

We may now assume that M_0 is the unique non-binary matroid labeling a vertex of T . Suppose there is a vertex labeled by a 3-connected binary matroid M_2 with at least four elements. Then, by Lemma 2.10, we see that M has an $M_0 \oplus_2 M_2$ -minor. Now M_0 has a $U_{2,4}$ -minor, and, as M_2 is 3-connected and binary, Lemma 3.2 tells us that M_2 has an $M(K_4)$ -minor. Let p_2 be the basepoint of $M_0 \oplus_2 M_2$. As above, M_0 has a $U_{2,4}$ -minor using p_2 . Since $\{M(K_4), U_{2,4}\}$ is 1-rounded [11], M_2 has an $M(K_4)$ -minor using p_2 . Thus M has a $U_{2,4} \oplus_2 M(K_4)$ -minor and therefore has a $U_{2,4} \oplus U_{1,1}$ -minor. Hence the lemma holds when M_2 exists.

We may now assume all the vertices in the tree decomposition other than the one labeled by M_0 are labeled by circuits or cocircuits. Suppose we have a path in our tree decomposition beginning at M_0 that has the form M_0 —circuit—cocircuit. Then M has a $U_{2,4} \oplus U_{0,1}$ -minor. By duality, if we have a path of the form M_0 —cocircuit—circuit, then M has a $U_{2,4} \oplus U_{1,1}$ -minor. As the lemma holds in these cases, we may assume the only non-trivial paths beginning at M_0 in the tree decomposition are of the form M_0 —circuit, or M_0 —cocircuit. In other words, M is obtained from M_0 by parallel extension of a subset of $E(M_0)$ and series extension of a disjoint subset of $E(M_0)$. $\square$

Recall that the matroid K is the matroid obtained from $U_{2,4}$ by adding elements in parallel to three of its elements.

Lemma 3.6. *The matroid R_6 is the only connected, but not 3-connected, member of $EX(\mathcal{Z})$. The connected, but not 3-connected, members of $EX(\mathcal{R})$ are R_6 , K , and K^* .*

Proof. Suppose $\mathcal{U} \in \{\mathcal{Z}, \mathcal{R}\}$. By Lemmas 2.4 and 2.7, R_6 is in $EX(\mathcal{U})$ and K and K^* are in $EX(\mathcal{R})$. Let M be a 2-connected member of $EX(\mathcal{U})$ that is not 3-connected and is not isomorphic to R_6 , K , or K^* . By Lemma 3.5, M

is obtained from a 3-connected non-binary matroid M_0 by parallel extension of a subset T of $E(M_0)$ and series extension of a disjoint subset S of $E(M_0)$ where $S \cup T \neq \emptyset$.

If $M_0 \cong U_{2,4}$, we will consider the cases $\mathcal{U}=\mathcal{Z}$ and $\mathcal{U}=\mathcal{R}$ separately. If $\mathcal{U}=\mathcal{Z}$, then M is in $\mathcal{U}$, as it satisfies (ii)(d) in Theorem 2.3, a contradiction. Now let $\mathcal{U}=\mathcal{R}$. As M has neither K nor K^* as a minor, both S and T have size less than three. By duality, we may assume that $0 \leq |T| \leq |S| \leq 2$. In each case, M can be realized as a relaxation of a binary matroid. For example, when $|S| = |T| = 2$, we can obtain M by relaxing the circuit-hyperplane in $M(G)$ where G is obtained from $K_{2,3}$ by adding edges in parallel to each edge meeting some fixed degree-2 vertex. The other cases can be checked similarly. We deduce that $M \in \mathcal{U}$, a contradiction.

We may now assume $|E(M_0)| \geq 5$, and consider $\mathcal{U} \in \{\mathcal{Z}, \mathcal{R}\}$. By switching to the dual if necessary, we may also assume that M has at least one non-trivial parallel class. Let $\{x, y\}$ be in that class.

3.6.1. *$M \setminus x$ can be obtained from a binary matroid by relaxing a circuit-hyperplane.*

Proof. We know $M \setminus x$ is in $\mathcal{U}$. Thus it satisfies one of (ii)(b)-(ii)(d) in Theorem 2.3. Assume $M \setminus x$ satisfies (ii)(b). Then $M \setminus x$ is isomorphic to a parallel extension of $U_{2,n}$, for some $n \geq 5$. Hence M is also a parallel extension of this matroid. Thus $M \in \mathcal{Z}$ so we may assume $\mathcal{U}=\mathcal{R}$. But M has a $U_{2,5}$ -minor, a contradiction to Lemma 2.7. Next assume that $M \setminus x$ satisfies (ii)(c). Then $M \setminus x$ is isomorphic to a series extension of $U_{n-2,n}$ for some $n \geq 5$, and M is a parallel extension of this series extension. Then $M \setminus x$, and hence M , has a $U_{2,4} \oplus U_{0,1}$ -minor, a contradiction. Lastly, assume $M \setminus x$ satisfies (ii)(d). Then $M \setminus x$ is obtained from $U_{2,4}$ by series extension of a subset S of $E(U_{2,4})$ and parallel extension of a disjoint subset T of $E(U_{2,4})$. Let $U = E(U_{2,4}) - S - T$. Recall that $\{x, y\}$ is a circuit of M . If y is in a non-trivial series class of $M \setminus x$, then M has a $U_{2,4} \oplus U_{0,1}$ -minor, a contradiction. Thus $y \in T \cup U$. Therefore M satisfies (ii)(d), so $M \in \mathcal{Z}$. Hence we assume $\mathcal{U}=\mathcal{R}$. As M has neither K nor K^* as a minor, $|S| < 3$ and $|T \cup y| < 3$. As noted above, in these cases, M can be realized as a relaxation of a binary matroid. $\square$

It is not hard to check that if $r(M_0) = 2$, then either $M \in \mathcal{U}$, or M contains an excluded minor. Thus we may assume $r(M_0) \geq 3$, and, by duality, $r^*(M_0) \geq 3$. As M_0 is non-binary and 3-connected, and all of P_6 , Q_6 , and $U_{3,6}$ are either in $EX(\mathcal{U})$ or contain members of $EX(\mathcal{U})$, Theorem 3.2 implies that M_0 contains a $\mathcal{W}^3$ -minor. By 3.6.1, $M \setminus x$ is a relaxation of a binary matroid N . Thus N has a circuit-hyperplane B such that relaxing B produces $M \setminus x$. If $y \notin B$ and N_1 is obtained from N by adding x back in parallel to y , then B is a circuit-hyperplane of N_1 whose relaxation is M , a contradiction.

We may now assume $y \in B$. Recall M_0 has a $\mathcal{W}^3$ -minor, and no P_6 -, Q_6 -, or $U_{3,6}$ -minor. As $\{\mathcal{W}^3, P_6, Q_6, U_{3,6}\}$ is 1-rounded [5], M_0 has a $\mathcal{W}^3$ -minor M_y using y . By Lemma 3.4, we know that y is a rim element of M_y . This implies that M has a $\mathcal{W}^3$ -minor in which one of the rim elements is replaced by the parallel class containing $\{x, y\}$. This is a contradiction since it implies M has a $U_{2,4} \oplus U_{0,1}$ -minor. $\square$

In finding the complete list of 3-connected excluded minors for each class, we will use the following lemmas, the second of which is from [7].

Lemma 3.7. *Let a matroid M be obtained from a matroid N by relaxing a circuit-hyperplane C . If M is 3-connected, then either N is 3-connected, or N has a unique 2-separation $(C, E(N) - C)$.*

Proof. Assume N has a 2-separation (X, Y) that is not $(C, E(N) - C)$. As M is obtained by relaxing C in N , for every subset S of $E(N)$ other than C , we have $r_N(S) = r_M(S)$. Thus $r_N(X) = r_M(X)$ and $r_N(Y) = r_M(Y)$. As relaxing a circuit-hyperplane does not change the rank of the matroid, (X, Y) is a 2-separation in M , a contradiction. $\square$

Lemma 3.8. *Let M be a non-binary matroid such that, for some element e , both M/e and $M \setminus e$ are binary. Then M is obtained from a 4-point line having ground set $\{e, e_1, e_2, e_3\}$ by a sequence of at most three 2-sums where the basepoints of these 2-sums are e_1, e_2 , and e_3 , the other part of each 2-sum is binary, and each of e_1, e_2 , and e_3 is the basepoint of at most one of these 2-sums.*

Lemma 3.9. *Let M be a matroid obtained from $U_{2,n}$ or $U_{n-2,n}$, for some $n \geq 4$, by a single-element extension or a single-element coextension. Then either M does not have a $\mathcal{W}^3$ -minor, or M has a Q_6 -minor.*

Proof. Assume M is obtained from $U_{2,n}$ by a single-element extension. Then M has no $\mathcal{W}^3$ -minor, as $r(M) = 2$. Now assume M is obtained from $U_{2,n}$ via coextension by a single element y . Assume M has a $\mathcal{W}^3$ -minor N . As $U_{2,n}$ has no parallel elements, y is not in any dependent rank-2 flats of M . Thus M has a free extension of $\mathcal{W}^3$ as a minor. Hence M has a Q_6 -minor. The result follows by duality. $\square$

Lemma 3.10. *Let the matroid N' be obtained from the matroid N by relaxing a circuit-hyperplane X of the latter. If N is non-binary, then so is N' .*

Proof. Assume N' is binary. Then, by Lemma 2.5, N must have at least two connected components including at least one, say N_1 , that is non-binary. As $r(N_1) \geq 2$, and X is a circuit-hyperplane, we know $X \subseteq E(N_1)$. Thus N has no loops, so $N = N_1 \oplus N_2$ where $r(N_2) = 1$. Let $\{e\}$ be a basis of N_2 . Let $N_1/I \setminus I^* \cong U_{2,4}$, where I and I^* are independent and coindependent in

N . Then I is also independent in N' . Thus, $N'/I/e$ has rank two with at least four rank-one flats, so N' has a $U_{2,4}$ -minor, a contradiction. $\square$

Lemma 3.11. *If a matroid N' is 3-connected and is obtained from a non-binary matroid N by relaxing a circuit-hyperplane X , then*

- (i) N' has a $U_{3,6}$ -, P_6 -, Q_6 -, R_6 -, $U_{2,4} \oplus U_{0,1}$ -, or $U_{2,4} \oplus U_{1,1}$ -minor; or
- (ii) N' has a matroid in the class $\mathcal{D}$ as a minor; or
- (iii) N' is isomorphic to $U_{2,n}$ or $U_{n-2,n}$ for some $n \geq 5$.

Proof. As N is non-binary and has a circuit-hyperplane, $|E(N)| \geq 5$. If $|E(N)| = 5$, then either N is $U_{2,4} \oplus U_{1,3}$, in which case N' is $U_{2,5}$, or N is $U_{2,4} \oplus U_{2,3}$, in which case N' is $U_{3,5}$. Now assume that the result holds for $|E(N)| = k$, and consider the case where $|E(N)| = k + 1 \geq 6$.

Suppose N has an element e such that $N \setminus e$ and N/e are both binary. Then, by Lemma 3.8, N is obtained from $U_{2,4}$, where $E(U_{2,4}) = \{e, e_1, e_2, e_3\}$, by a sequence of at most three 2-sums where the basepoints are e_1, e_2 , and e_3 , the other part of each 2-sum is binary, and each of e_1, e_2 , and e_3 is the basepoint of at most one 2-sum. As N' is 3-connected, by Lemma 3.7, N has only one 2-separation, and the circuit-hyperplane X must be one part of this 2-separation. Without loss of generality, assume that the basepoint of the corresponding 2-sum is e_3 , and let L denote the binary matroid being 2-summed with $U_{2,4}$. Assume $X = \{e, e_1, e_2\}$. Then $r(\{e, e_1, e_2\}) = r(N) - 1 = 2$, so $r(L) = 2$. As N is non-binary, L must have at least three rank-one flats. But, as L is binary, it cannot have four such flats. Since $|E(N)| \geq 6$, the matroid L is a parallel extension of $U_{2,3}$. Therefore N' is a parallel extension of $U_{3,5}$, so N' contains a $U_{2,4} \oplus U_{0,1}$ -minor.

We may now assume that $X = L \setminus e_3$. Consider N^* . The set $\{e, e_1, e_2\}$ is a circuit-hyperplane in N^* whose relaxation produces $(N')^*$. Thus, by the argument above, $(N')^*$ has a $U_{2,4} \oplus U_{0,1}$ -minor. Hence N' has a $U_{2,4} \oplus U_{1,1}$ -minor. Therefore we may assume that, for every $e \in E(N)$, at least one of $N \setminus e$ and N/e is non-binary.

Assume N/e is non-binary for some $e \in X$. By Lemma 2.2, N'/e is obtained from N/e by relaxing $X - e$. Suppose N'/e is 3-connected. Then, by our induction hypothesis, either N'/e has one of the specified minors, so N' also does; or N'/e is isomorphic to $U_{2,n}$ or $U_{n-2,n}$. In the latter case, as N' is 3-connected, either $N' \cong U_{n-1,n+1}$ and the lemma holds, or both the rank and corank of N' are at least three. As N' is non-binary, Theorem 3.2 implies that N' must contain a $\mathcal{W}^3$ -, Q_6 -, P_6 -, or $U_{3,6}$ -minor. By Lemma 3.9, the only way we can coextend $U_{2,n}$ to get a matroid with a $\mathcal{W}^3$ -minor gives us a matroid that has a Q_6 -minor, so we are done. Thus we may assume that N'/e is not 3-connected. As N' is 3-connected, N'/e must be connected. Furthermore, by Lemma 3.10, since N/e is non-binary, so is N'/e .

By Lemma 3.5, either N'/e has an R_6 -, $U_{2,4} \oplus U_{0,1}$ -, or $U_{2,4} \oplus U_{1,1}$ -minor, or N'/e can be obtained from a 3-connected non-binary matroid M_0 by parallel extension of a subset of $E(M_0)$, and series extension of a disjoint

subset of $E(M_0)$. In the first case, the lemma holds. Thus we can assume the second case holds. As N' is 3-connected, N'/e cannot have any elements in series. Since N'/e is not 3-connected, it must have parallel elements. By the 1-roundedness of $\{U_{2,4}\}$, we know N'/e has a $U_{2,4} \oplus_2 U_{1,3}$ -minor. As N' is 3-connected, when we coextend N'/e by e , we must destroy the non-trivial parallel classes in N'/e . Thus N' has at least one dependent flat of rank two. It is not hard to check that each coextension of $U_{2,4} \oplus_2 U_{1,3}$ that has no parallel elements has a $U_{2,4} \oplus U_{1,1}$ -, Q_6 -, P_6 -, or R_6 -minor.

By duality, the required result holds if $N \setminus f$ is non-binary for some f in $E(N) - X$. We may now assume, since N has no element z such that $N \setminus z$ and N/z are both binary, that $X = \{e : N/e \text{ is binary}\}$ and $E(N) - X = \{f : N \setminus f \text{ is binary}\}$. By Theorem 2.3, we deduce that one of (ii)(a)-(ii)(d) holds.

If N satisfies (ii)(b), then one parallel class must have size two to ensure N has a circuit-hyperplane. The other parallel classes must be trivial, otherwise N' would not be 3-connected. Thus $N' \cong U_{2,n+1}$, where N is a parallel extension of $U_{2,n}$. By duality, if N satisfies (ii)(c), then $N' \cong U_{n-1,n+1}$. If N satisfies (ii)(a), then N can be obtained from some binary matroid M by relaxing a circuit-hyperplane Y in M . This gives us two partitions of the ground set, $(X, E(N) - X)$ and $(Y, E(N) - Y)$. For all $e \in Y$, we know $N \setminus e$ is binary, by Lemma 2.2. Therefore $Y \subseteq E(N) - X$. Similarly, for all $e \in E(N) - Y$, the matroid N/e is binary, hence $E(N) - Y \subseteq X$. Thus $X = E(N) - Y$ and $E(N) - X = Y$. Hence N' is obtained from the connected binary matroid M by relaxing the two disjoint circuit-hyperplanes X and Y . Therefore N' is in $\mathcal{D}$.

Finally, assume N satisfies (ii)(d). Then N can be obtained from $U_{2,4}$ by series extension of a subset S of $E(U_{2,4})$ and parallel extension of a disjoint subset T of $E(U_{2,4})$, where S or T may be empty. As N' is 3-connected, $|S| + |T| \leq 1$ by Lemma 3.7. It follows easily that $|E(N')| \leq 5$; a contradiction. $\square$

Lemma 3.12. *The complete list of 3-connected members of $EX(\mathcal{Z})$ is Q_6 , P_6 , $U_{3,6}$, and the matroids in $\mathcal{D}$. The complete list of 3-connected members of $EX(\mathcal{R})$ is $U_{2,5}$, $U_{3,5}$, and the matroids in $\mathcal{D}$*

Proof. Suppose $\mathcal{U} \in \{\mathcal{Z}, \mathcal{R}\}$. Let M be a 3-connected excluded minor of $\mathcal{U}$ that is not isomorphic to Q_6 , P_6 , $U_{3,6}$, $U_{2,5}$, $U_{3,5}$, or any of the matroids in $\mathcal{D}$. Either M is a relaxation of a non-binary matroid, or M is not a relaxation of any matroid at all. In the first case, by Lemma 3.11, M has a minor isomorphic to Q_6 , P_6 , $U_{3,6}$, R_6 , $U_{2,4} \oplus U_{0,1}$, $U_{2,4} \oplus U_{1,1}$, or some matroid in $\mathcal{D}$; or M is isomorphic to $U_{2,n}$ or $U_{n-2,n}$ for some $n \geq 5$. All of these provide contradictions, so we may assume that M cannot be realized as a relaxation.

Clearly we may assume $r(M) \geq 3$ and $r^*(M) \geq 3$. Thus M must contain one of $\mathcal{W}^3$, Q_6 , P_6 , and $U_{3,6}$ as a minor, by Theorem 3.2. As all of these

except $\mathcal{W}^3$ contain excluded minors of $\mathcal{U}$, we know that M has a $\mathcal{W}^3$ -minor. Let $\mathcal{W}^k$ be the largest whirl-minor of M . By Seymour's Splitter Theorem, we may grow from $\mathcal{W}^k$ by single-element extensions and single-element co-extensions, staying 3-connected, until we reach M . Let x be the last element added in this way. By duality, we may assume that x is added via extension. Thus $M \setminus x$ is a non-binary 3-connected member of $\mathcal{U}$. If $\mathcal{U} = \mathcal{R}$, then $M \setminus x$ is a relaxation of a binary matroid. If $\mathcal{U} = \mathcal{Z}$, then $M \setminus x$ satisfies one of (ii)(a)-(ii)(d) in Theorem 2.3. As $M \setminus x$ is 3-connected, it cannot satisfy (ii)(b)-(ii)(d). Hence, in both cases, $M \setminus x$ is a relaxation of a binary matroid N_1 .

Let B be the special basis in $M \setminus x$ that is a circuit-hyperplane in N_1 . For all $e \in E(M \setminus x) - B$, the set $B \cup e$ is a circuit in $M \setminus x$. Now B is also a basis of M , and $B \cup e$ is a circuit of M for all $e \in E(M) - (B \cup x)$. If $B \cup x$ is a circuit of M , then M can be realized as a relaxation of some matroid by Lemma 3.3, a contradiction. Thus $C_M(x, B) \neq B \cup x$. Therefore there is some $y \in B$ such that $y \notin C_M(x, B)$. Now, by Lemma 2.2, $M \setminus x / y$ can be obtained by relaxing the circuit-hyperplane $B - y$ in $N \setminus x / y$.

Assume that $M \setminus x / y$ is binary. As $M \setminus x / y$ is obtained by relaxing the circuit-hyperplane $B - y$ in the binary matroid $N \setminus x / y$, it follows that $M \setminus x / y$ can be obtained from a circuit C by adding some, possibly empty, set of elements in parallel with some element z of C where $C - z = B - y$ (see, for example, [9, p.386]). As $M \setminus x$ is 3-connected, it has no non-trivial series classes. Hence $|C - z| = 1$, so $r(M \setminus x / y) = 1$; a contradiction. Therefore we know that $M \setminus x / y$ must be non-binary. Thus M / y is also non-binary. Here we consider the cases $\mathcal{U} = \mathcal{Z}$ and $\mathcal{U} = \mathcal{R}$ separately. Our goal in each case is to show the following.

3.12.1. *M / y can be obtained from a binary matroid via relaxation.*

Proof. This is certainly true if $\mathcal{U} = \mathcal{R}$. Now assume $\mathcal{U} = \mathcal{Z}$. Then M / y satisfies one of (ii)(a)-(ii)(d) in Theorem 2.3. First note that M / y cannot satisfy (ii)(c), because a connected single-element coextension of a series extension of $U_{n-2, n}$ has corank two, and so has no $\mathcal{W}^3$ -minor. Assume M / y satisfies (ii)(b). Then M / y is a parallel extension of $U_{2, n}$ for some $n \geq 5$, and $r(M) = 3$. As M has a $\mathcal{W}^3$ -minor, it is routine to check that we must coextend M / y by y in a way that creates a matroid having a minor isomorphic to $U_{2, 4} \oplus U_{1, 1}$, Q_6 , or P_6 . We may now assume M / y satisfies (ii)(d). As M is 3-connected, M cannot have any non-trivial series classes. It is routine to check that either M contains an excluded minor, or M can be realized as a relaxation of a binary matroid. $\square$

We now revert to working in generality, where $\mathcal{U} \in \{\mathcal{Z}, \mathcal{R}\}$. We know that $M \setminus x$ is obtained by relaxing a circuit-hyperplane B in a binary matroid N_1 , and M / y is obtained by relaxing a circuit-hyperplane B' in a binary matroid N_2 , where $y \in B$. We show next that

3.12.2. *$B' = B - y$.*

Proof. By Lemma 2.2, $M \setminus x/y$ is obtained by relaxing the circuit-hyperplane $B - y$ in N_1/y . Consider $(M \setminus x/y) \setminus e$ for $e \notin B - y$. Assume $(M \setminus x/y) \setminus e$ is binary. Then, as $(M \setminus x/y) \setminus e$ is a relaxation of a binary matroid, we know $(M \setminus x/y) \setminus e \cong U_{t-1,t} \oplus_2 U_{1,v}$ for some $t, v \geq 2$. Now $E(M \setminus x/y \setminus e)$ has a partition (S, P) where S is the relaxed set $B - y$ and P is its complement. Then $S \cup y$ is the relaxed set of $M \setminus x$, and $P \cup e$ is the relaxed set of $(M \setminus x)^*$. As $r_{M \setminus x}(P \cup y) = 2$, we know $|P| \leq 2$. Otherwise the matroid N_1 would be non-binary. Similarly, $r_{M \setminus x}^*(S \cup e) = 2$. Thus $|S| \leq 2$. However, $r(M \setminus x) = |S \cup y|$, and $r^*(M \setminus x) = |P \cup e|$. Thus, as $M \setminus x$ has a $\mathcal{W}^k$ -minor for some $k \geq 3$, we know $|S \cup y| = |P \cup e| = 3$. Therefore, $M \setminus x$ has six elements, so $M \setminus x \cong \mathcal{W}^3$. It is not hard to show that the only 3-connected single-element extension of $\mathcal{W}^3$ that does not contain an excluded minor is the non-Fano matroid F_7^- , depicted in Figure 3. Thus M must be F_7^- . But F_7^- is a relaxation of the Fano plane, which is binary, a contradiction. Thus we may assume $(M \setminus x/y) \setminus e$ is non-binary for all $e \notin B - y$. Then, for all such e , the matroid $M/y \setminus e$ is also non-binary. By Lemma 2.2, for every $e \in B'$, the matroid $M/y \setminus e$ is binary. Thus if e is not in $B - y$, then it is not in B' . Therefore, $B' \subseteq B - y$. As $B - y$ and B' are both bases for M/y , the result holds. $\square$

We have $C_M(e, B) = B \cup e$ for all $e \in E(M) - (B \cup x)$. We also know $C_{M/y}(e, B - y) = (B - y) \cup e$ for all $e \in E(M) - B$. Thus, since $C_M(x, B) \neq B \cup x$, we have $C_M(x, B) = (B - y) \cup x$. We know that $M \setminus x$ has the basis B that results from relaxing a circuit-hyperplane in N_1 . As $M \setminus x$ has a $\mathcal{W}^3$ -minor, but no Q_6 -, P_6 , or $U_{3,6}$ -minor, the 1-roundedness of $\{\mathcal{W}^3, Q_6, P_6, U_{3,6}\}$ implies that $M \setminus x$ has a $\mathcal{W}^3$ -minor using y . By Lemma 3.4, as y is in B , it follows that this $\mathcal{W}^3$ -minor has y as a rim element. Hence by adding x back, M has a single-element extension of $\mathcal{W}^3$ as a minor. Let $\{b_1, b_2, y\}$ be the set of rim elements in $\mathcal{W}^3$. There are only two single-element extensions of $\mathcal{W}^3$ that do not contain excluded minors. These are F_7^- and a parallel extension of a spoke element of $\mathcal{W}^3$ (see Figure 3). But, in each of them, the set $\{b_1, b_2, x\}$ should be a circuit because $(B - y) \cup x$ is a circuit of M and the only elements of $M \setminus x$ that can be contracted to produce the $\mathcal{W}^3$ -minor must belong to B . This contradiction completes the proof. $\square$

Proofs of Theorems 1.1 and 1.2. These follow immediately by combining Lemmas 3.1, 3.6, and 3.12. $\square$

4. THE COMPLEXITY OF $\mathcal{D}$

Jim Geelen asked (private communication) whether members of $\mathcal{D}$ could contain arbitrarily large projective geometries. In this section, we observe that they can. Note that all sums in this section are modulo two. Let A be a $k \times (2^k - 1)$ matrix representing the rank- k binary projective geometry




FIGURE 3. Geometric representations of F_7^- , and a parallel extension of a spoke element of $\mathcal{W}^3$.

$$\begin{array}{c}
 \begin{array}{cccc} 1 & 2 & 3 & \dots \end{array} \\
 \left[\begin{array}{c|c} I_{n-1} & \begin{array}{c} 1 \ 0 \\ 1 \ 0 \\ \vdots \ \vdots \\ 1 \ 0 \\ 1 \ 1 \end{array} \end{array} \right] \begin{array}{c} n \ n+1 \end{array} \dots \begin{array}{c} 2n \end{array} \\
 \begin{array}{cccc} 0 & 0 & 0 & \dots \end{array} \end{array} \left[\begin{array}{c|c|c} \begin{array}{c} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_t \end{array} & \begin{array}{c} A \\ I_k \end{array} & \begin{array}{c} I_k \\ A^T \end{array} \end{array} \right] \begin{array}{c} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_t \end{array} \\
 \begin{array}{cccc} 0 & 0 & 1 & \dots \end{array} \left[\begin{array}{c|c|c} \begin{array}{c} \alpha_1+1 \ \alpha_2+1 \end{array} & \begin{array}{c} \alpha_t+1 \end{array} & \begin{array}{c} \gamma \end{array} \end{array} \right] \begin{array}{c} \gamma \end{array} \\
 \begin{array}{cccc} 1 & 1 & \dots & 1 \end{array} \left[\begin{array}{c|c|c} \begin{array}{c} 1 \end{array} & \begin{array}{c} 1 \end{array} & \begin{array}{c} 1 \end{array} \end{array} \right] \begin{array}{c} 1 \end{array}
 \end{array}$$

FIGURE 4. The matrix Z .

$PG(k-1, 2)$, where k is odd. Let $n = 2^k + k + 1$, let $t = 2^k + k - 1$, and consider the rank- n binary matrix Z in Figure 4.

Let z_{sc} denote the entry in row s and column c of Z . Let $\alpha_i = \sum_{s=1}^{n-2} z_{s(n+1+i)}$, for $1 \leq i \leq t$. Let $\beta_j = \sum_{c=n+2}^{2n-1} z_{jc}$, for $1 \leq j \leq t$, and let $\gamma = 1 + \sum_{j=1}^t \beta_j$. Let Z' be the submatrix of Z whose columns are labeled by $n+1, n+2, \dots, 2n$. Then each column in Z' is contained in the hyperplane of $PG(n-1, 2)$ consisting of those vectors whose coordinates sum to zero. Moreover, no other column of Z is in this hyperplane. The definitions ensure that all the rows of Z' , except possibly row $n-1$, sum to zero. To see that row $n-1$ also sums to zero, note that $\sum_{j=1}^t \beta_j = \sum_{i=1}^t \alpha_i$ since both of these sums count the number of non-zero entries in the same submatrix. We know that $\sum_{c=n+2}^{2n-1} z_{(n-1)c} = 1 + \sum_{i=1}^t (\alpha_i + 1) = 1 + t + \sum_{i=1}^t \alpha_i$. As t is even, $1 + t + \sum_{i=1}^t \alpha_i = 1 + \sum_{i=1}^t \alpha_i = 1 + \sum_{j=1}^t \beta_j = \gamma$. Thus $\{n+1, n+2, \dots, 2n\}$ is a circuit-hyperplane of $M[Z]$, and it is easy to see that its complement is as well. By relaxing both these circuit-hyperplanes, we get a member of $\mathcal{D}$ that contains a $PG(k-1, 2)$ -minor.

REFERENCES

- [1] R.E. Bixby, l -matrices and a characterization of non-binary matroids, *Discrete Math.* 8 (1974) 139-145.
- [2] W.H. Cunningham, A combinatorial decomposition theory, Ph.D. thesis, University of Waterloo (1973).
- [3] J. Kahn, A problem of P. Seymour on nonbinary matroids, *Combinatorica* 5 (1985) 319-323.
- [4] D. Mayhew, G. Whittle, and S. van Zwam, Stability, fragility, and Rota's Conjecture, *J. Combin. Theory Ser. B*, to appear.
- [5] J.G. Oxley, On the intersections of circuits and cocircuits in matroids, *Combinatorica* 4 (1984) 187-195.
- [6] J.G. Oxley, A characterization of certain excluded-minor classes of matroids, *European J. Combin.* 10 (1989) 275-279.
- [7] J.G. Oxley, A characterization of a class of non-binary matroids, *J. Combin. Theory Ser. B* 49 (1990) 181-189.
- [8] J. Oxley, *Matroid Theory*, Second edition, Oxford University Press, New York, 2011.
- [9] J. Oxley and G. Whittle, On weak maps of ternary matroids, *European J. Combin.* 19 (1998) 377-389.
- [10] P.D. Seymour, Decomposition of regular matroids, *J. Combin. Theory Ser. B* 28 (1980) 305-359.
- [11] P.D. Seymour, Minors of 3-connected matroids, *European J. Combin.* 6 (1985) 375-382.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA, USA

E-mail address: `oxley@math.lsu.edu`

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA, USA

E-mail address: `JTAYL75@math.lsu.edu`